

Potential Physiological Importance of Pyrroloquinoline Quinone

Robert Rucker, PhD; Winyoo Chohanadisai, PhD; Masahiko Nakano, PhD

Abstract

Pyrroloquinoline quinone (PQQ) is a novel biofactor for which a proposition can be made for physiological importance. PQQ was first recognized as an enzyme cofactor in bacteria. It has recently been tentatively identified as a component of interstellar dust. Thus, PQQ may have been present throughout early biological conception and evolution. PQQ is also a potent plant growth factor. Consequently, for animals and humans, there has been constant exposure to PQQ. In animals, PQQ is reported to participate in a range of biological functions with apparent survival benefits (e.g., improved neonatal growth and reproductive performance). There are also benefits from PQQ supplementation related to cognitive, immune, and antioxidant functions, as well as protection from cardiac and neurological ischemic events. Although PQQ is not currently viewed as a vitamin, its involvement in cell signaling pathways, particularly those important to mitochondriogenesis in experimental animal models, may eventually provide a rationale for defining PQQ as vital to life. For humans, such evidence suggests there may be similar parallels or benefits from improving PQQ status. (*Altern Med Rev* 2009;14(3):268-277)

Introduction

Pyrroloquinoline quinone (PQQ) was first recognized as a bacterial cofactor by Hauge,¹ and later by Anthony,²⁻⁴ Salisbury,^{5,6} Duine,⁷ and their co-workers. PQQ, also known as methoxatin (Figure 1), is water soluble and heat stable. Under appropriate conditions, PQQ is capable of catalyzing continuous redox cycling (the ability to catalyze repeated oxidation and reduction reactions), as well as oxidative deaminations.⁸

These chemical properties are novel in many respects. For example, in chemical assays, PQQ's stability renders it capable of carrying out thousands of redox catalytic cycles; whereas, other bioactive quinones capable of redox cycling (e.g., epicatechin) tend to self oxidize and/or form polymers (e.g., tannins). Table 1 contains data that in part demonstrates the effectiveness of PQQ as a redox cycling agent.⁸⁻¹³ There is also a range of papers that describe PQQ's use in an analytical setting.¹⁴⁻¹⁷ PQQ molecules can be immobilized and fixed at the surface of analytical electrodes. When coupled to appropriate enzyme systems, highly specific and sensitive assays have been developed to assay compounds ranging from glucose to common narcotics.¹⁴⁻¹⁷ As will be highlighted in subsequent sections, the novel chemical attributes of PQQ help explain many of its metabolic and health-related features.


Biochemical Roles and Mechanisms Evolution and Its Functions in Bacteria and Plants

To make the case for physiological and biomedical importance in humans, it is important to note that many of PQQ's functions are universal. For example, for many bacterial species, PQQ stimulates growth and serves as a cofactor for a special class of dehydrogenases/

Robert Rucker, PhD – Professor, Nutrition Department, UC Davis, Davis, CA.
Correspondence address: UC Davis, Nutrition Department, One Shields Avenue, Davis, CA 95616.
Email: rbrucker@ucdavis.edu

Winyoo Chohanadisai, PhD – Post-graduate Research Fellow, Nutrition Department, UC Davis, Davis, CA.

Masahiko Nakano, PhD – Program Officer and Research Director, Developmental Analysis Section, Niigata Research Laboratory, Mitsubishi Gas Chemical Company, Inc., Tayuhama, Niigata, Japan.

Figure 1. Structure of PQQ

oxidoreductases. Enzymes containing PQQ are sometimes designated quinoproteins.⁸ Although the “quinoproteins” include many types of quinone-containing proteins and enzymes, the PQQ-requiring glucose and alcohol dehydrogenases are distinguished, because the PQQ associated with these proteins is dissociable and synthesized in metabolic pathways that can be separately controlled from those pathways important for the generation of the eventual targeted protein.¹⁸⁻²⁰ In addition to a cofactor role, PQQ can also be thought of as a trophic factor important to the growth and metabolism of bacteria, particularly methylotrophic bacteria (bacteria capable of growing on simple carbon sources).

From an evolutionary perspective, current evidence suggests PQQ is a component of interstellar dust as analyzed by particle impact time-of-flight mass spectrometry.^{21,22} Cometary grains are considered to be the precursors of organic materials in early life on the earth. It can also be argued that strong redox catalysts would be required to trigger the earliest chemical evolutionary steps. The presence of PQQ in stellar dust raises the question of PQQ's evolutionary importance to simpler life forms, given its wide range of chemical properties, such as redox catalysis and the ability to carry out useful amino acid modifications (e.g., oxidative deamination reactions).

At the next level it is important to highlight the symbiotic relationship between plants and soil bacteria, such as rhizobacterium.²³⁻²⁵ Plants cultivated in hydroponic culture systems with rhizobacterium have significantly increased height, flower number, fruit number, and total fruit weight; whereas, this does not occur with genetically modified rhizobacteria unable to produce PQQ.²³ PQQ added directly to hydroponic culture systems also confers a significant increase in the fresh weight of seedling plants. In part, the role of PQQ is related to phosphate uptake by plants, because PQQ, as a cofactor for rhizobacteria dehydrogenases, facilitates making soil and the local environment more acidic.²⁴ As a consequence, phosphate is made more available to plant roots. In addition, independent roles have been proposed for PQQ related to plant growth via activation of cell signaling, antioxidant defense, and viral protection.²⁵

For humans and animals, the ubiquitous presence of PQQ in common types of bacteria, soil, and plants suggests constant exposure to PQQ. PQQ has been found in all plant foods analyzed to date.^{8,26-28} In this regard, it is interesting to note that although many bacteria make PQQ, this is not the case for common intestinal bacteria, such as *Escherichia coli*. *Escherichia coli* can synthesize PQQ-dependent enzymes capable of utilizing PQQ under certain nutrient limiting conditions;²⁹ however, the enzymes only become functional when PQQ is present. Hence, an external source of PQQ may be important in sustaining human and animal tissue levels of PQQ, as well as maintaining an optimal enteric environment.

Table 1. PQQ as a Redox Cycling Agent

Compound	Potential Number of Catalytic Cycles
PQQ	20,000
Quercetin	800
Catechin	75
Epicatechin	700
Norepinephrine	200
Epinephrine	100
DOPA	20
6-OH-DOPA	20
Ascorbic Acid	4

Redox cycling systems result in repeated chemical reactions in which molecules that act as catalysts are repeatedly oxidized and/or reduced. The potential number of catalytic cycles (number of repeated reactions) depends in part on chemical stability. PQQ is relatively stable; whereas, self-oxidation, polymerization, and/or changes in chemical structure are factors that compromise the chemical stability of many bioactive quinones or enediols (e.g., ascorbic acid). Details of the redox cycling system and basis for defining the relative number of catalytic cycles may be found in references 8, 13 and 67.

Mechanisms and Proposed Functions in Humans and Animals

A number of physiological properties have been attributed to PQQ, ranging from classical water-soluble vitamin/cofactor functions to those important to antioxidant potential.^{8,30-39} While a role as a vitamin in animal or human nutrition seems unlikely at this time, similar to other polyphenolic biofactors, there is strong evidence PQQ may play an important role in pathways important to cell signaling.³⁵⁻³⁹ PQQ can also serve as an antioxidant.⁴⁰ The importance of PQQ to mammalian health is evident when it is omitted from chemically defined diets, resulting in a wide range of systemic responses, including growth impairment, compromised immune responsiveness, and abnormal reproductive performance in mouse and rat experimental models.^{8,25,26,41,42} Furthermore, varying PQQ in highly

refined diets causes modulation in mitochondrial content, alters lipid metabolism, and reverses inhibition elicited by classical complex I inhibitors.^{39,41-43}


Improvements in mitochondrial respiratory control are potentially important to a number of health issues, ranging from increased longevity to improved energy utilization and protection from reactive oxygen species. Mitochondrial DNA depletion and mutations are associated with cardiomyopathy, developmental delays, and impaired neurological and mitochondrial function,⁴⁴ which further highlights the importance of optimal mitochondrial function for health and well-being. Regarding possible mechanisms of PQQ action (Figure 2), given that many mitochondrial-related

events are regulated by peroxisome proliferator-activated receptor-gamma coactivator-1 alpha (PGC-1 α) and nuclear respiratory factors, an interaction between PQQ and a PGC-1 α -related pathway seems a logical possibility. Indeed, such interactions have recently been reported.⁴⁴⁻⁴⁶

PGC-1 α is a transcriptional coactivator that regulates genes involved in energy metabolism.^{45,46} An interaction with this protein and its association with multiple transcription factors can provide a direct link between an external physiological stimulus and the regulation of mitochondrial biogenesis. PGC-1 α is also a major factor that regulates muscle fiber type determination and appears to be involved in controlling blood pressure, regulating cellular cholesterol homeostasis, and the development of obesity. Moreover, PGC-1 α is associated with a reduction in reactive oxygen species and protection against various mitochondrial toxins.⁴⁵



Figure 2. Cell Signaling and PQQ



In addition to interacting with PGC-1α, PQQ can also affect the activity of ras,³⁸ an oncogene important to signal transduction processes involved in growth and development. PQQ can stimulate activated c-Ha-ras-transformed NIH3T3 mouse fibroblasts,³³ resulting in increased cell proliferation. With regard to ras, activation usually results in changes in cell growth, differentiation, and survival. In addition to ras,

Kumazawa et al³³ have also observed that activation of ERK occurs in response to adding PQQ to cultured fibroblasts. ERK is one of the many protein kinases that functions in the ras-signaling pathway to activate other components of transcription (e.g., activators, co-activators, and transcription factors). Similarly, raf is depicted in Figure 2, because like ERK it is also part of the chain of events that progress from ras to eventual

activation of signal transducers and activators of transcription (STAT) factors that are essential to control of cell growth, survival, and differentiation. Regarding PGC-1 α -related cell signaling,^{8,39,41-43} activation starts with signals from 5' AMP-activated protein kinase (AMPK) or one of the many mitogen-activated protein (MAP) kinases that are linked to various cell surface receptors.^{44,45} Such kinases activate cAMP response element binding protein (CREB), which is a transcription factor that binds to certain DNA sequences called cAMP response elements. The authors recently reported³⁹ that CREB activation in combination with other transcription factors, such as nuclear respiratory factors 1 and 2 (NRF) and mitochondrial transcription factors (e.g., Tfam), leads to increased mitochondrial biogenesis observed with PQQ administration.⁴¹⁻⁴³

PQQ has been reported to influence the activity of DJ-1³⁷. DJ-1 is involved in cellular oxidative stress responses, and autosomal-recessive mutations in DJ-1 lead to Parkinson's disease. A possible role of an interaction between DJ-1 and PQQ may be to facilitate and fine-tune overall cellular regulation.³⁷ The cell signals for growth and energy utilization via mitochondria are in communication with apoptotic (programmed cell death) signals. DJ-1, most likely by associating with Daxx (a multifunctional protein associated with apoptotic events), is capable of modulating apoptosis by inactivating yet another cell signaling pathway, the so-called Janus kinase (JNK) cell-signaling pathway. An important feature of this interaction is potential "crosstalk" with STAT components and control of caspase activation/deactivation (cysteine-aspartic acid proteases). Caspases are associated with apoptosis or in pathological situations with cellular necrosis and inflammation. Accordingly, when all is taken together, an important feature of the interactions depicted in Figure 2 is the crosstalk between the signals, such as ras, DJ-1, and numerous kinases, that control proliferation, apoptosis, and mitochondrial biogenesis.

Regarding PQQ's role as an antioxidant, recent studies on the aroxyl radical-scavenging action of reduced PQQH₂ have shown PQQ exists in cells in a reduced form.⁴⁰ Like vitamin C, glutathione, vitamin E, and uric acid, PQQ can act as an antioxidant.⁸ As examples, Tsuchida et al⁴⁷ and Urakami et al⁴⁸ reported PQQ protects against acute liver damage induced by agents

such as carbon tetrachloride or endotoxin. Hamagishi et al⁴⁹ observed that PQQ administered (i.p.) at 10 or 30 mg/kg body weight causes a decrease in carrageenan-induced edema by 39- and 76 percent, respectively. It is also noteworthy that on a molar basis, PQQ is a better inhibitor of tissue oxidation in peritoneal cells than α -tocopherol and ascorbic acid, following initiation by zymosan, carrageenan, or N-formyl-methionyl-leucyl-phenylalanine, all of which provoke inflammatory responses.⁴⁴ From a mechanistic perspective, in addition to serving as an antioxidant, the effects of PQQ on genes, such as PGC-1 α , DJ-1, and genes in the ras family help explain many of the physiological and clinical functions ascribed to PQQ.

Clinical Implications

The following subsections briefly describe clinical implications of PQQ use. Although much of this work was conducted in animal models, current efforts in humans and human cell lines demonstrate important parallels.

Improvements in Reproduction, Early Development, Growth, and Immune Function

Nutritional studies indicate PQQ can serve as a growth factor and improves neonatal survival.^{8,31,41,42} In human fibroblast cultures, PQQ enhances cell growth and proliferation when added to cell cultures.^{38,50} Signs of PQQ deprivation include friable skin, evidence of hemorrhage and diverticuli, and reduction in general fitness. The growth-related observations are novel in that adding 100-200 μ g PQQ/kg to purified diets improves growth, development, and reproductive parameters in rodent models.^{8,30} For perspective, the animal requirements for folic acid or for biotin range from 200-500 μ g/kg diet, respectively. These effects are similar to the improvements when more complex diets are fed (i.e., made of less refined ingredients).^{30,31} Moreover, in female mice and rats fed PQQ-deficient diets, fertility is decreased (fewer successful pregnancies and smaller litter size)^{30,31} compared to mice or rats fed PQQ-supplemented diets.

PQQ deprivation also results in defects in immune function and reduction in interleukin-2 (IL-2) levels. There is loss of B- and T-cell sensitivity to mitogens. The body normally produces IL-2 during an

immune response. IL-2 is necessary for the development of T-cell immunologic memory, one of the unique characteristics of the immune system. Maximizing sensitivity of B- and T-cells to mitogens is achieved in mice when as little as 1 nmol PQQ is added per gram of diet, about 100-400 µg PQQ per day in human equivalents.^{30,31}

PQQ and Neuroprotection

Neuronal cell death in experimental models of stroke and spinal cord injury is attenuated by PQQ.^{31,32,51-54} PQQ has been demonstrated to protect the redox modulatory site of N-methyl-d-aspartic acid (NMDA) receptors.^{36,51-59} Agents that protect NMDA-receptor function are often neuroprotective in experimental stroke and spinal cord injury models. In this regard, intraperitoneal administration of PQQ effectively promotes the functional recovery of spinal cord injury in rats after hemi-transection.³⁶ Protection is preceded by a decrease in inducible nitric oxide synthase (iNOS) mRNA. Nitric oxide is implicated in NMDA receptor-mediated neurotoxicity. Administration of PQQ decreased lesion size and increased axon density associated with the lesion area. Furthermore, recent studies suggest PQQ protects against secondary damage by reducing iNOS expression following a primary physical injury to the spinal cord. Peroxynitrite is a potential byproduct of abnormally high nitric oxide (NO) or cellular hydrogen peroxide levels. The demonstration that iNOS expression is reduced is in part a validation of previous work showing that PQQ treatment suppresses peroxynitrite formation.⁵⁶ Moreover, PQQ's ability to affect the oxidative status of DJ-1 adds an additional dimension.³⁷ As has been noted previously, the expression level and oxidation status of DJ-1 have been shown to play a role in antioxidative stress reactions important to neurological function. These findings add to the initial observations by Jensen et al⁵⁴ that PQQ effectively reduces infarct size in an experimental model of cerebral hypoxia/ischemia. PQQ administered i.p. at 10-15 mg/kg body weight in rats was effective in reducing cerebral infarct volumes measured 72 hours or more after a neurovascular insult. Three hours after ischemia a dose of 3 mg/kg significantly reduces infarct volume compared to vehicle-treated animals. These data indicate PQQ may be a useful neuroprotectant in stroke therapy.

Even at a more subtle level, PQQ exposure can affect learning ability and memory function in rats.⁶⁰ Rats fed a PQQ-supplemented diet demonstrate improved learning using the Morris water maze test as an index. Rats were fed 20 mg PQQ, 300 mg coenzyme Q10 (CoQ10), 200 mg R,R,R- α -tocopherol, or 20 mg PQQ + 300 mg CoQ10/kg body weight/day for nine weeks (from age four weeks). Each rat was subjected to hyperoxia as the oxidative stress (using a 100% oxygen chamber) for 48 hours. Those fed PQQ-supplemented diets were protected from a memory deficit that was apparent in controls not fed PQQ. As a novel control, rats fed vitamin E-supplemented and -deficient diets were tested. Vitamin E-deficient rats fed PQQ and/or CoQ10 demonstrated improved learning function. In addition, longer-term memory function was maintained independently by PQQ, but not by CoQ10 supplementation. Thus, PQQ seems potentially effective in sustaining learning functions during oxidative stress, independent of and in a manner different from that of vitamin E.

PQQ and Cardiac Function

PQQ is useful in models of cardiac ischemia.^{43,61,62} PQQ confers resistance to acute oxidative stress in freshly isolated cardiomyocytes.⁶¹ Both oxidative damage and mitochondrial membrane potential depolarization (induced by hydrogen peroxide) are significantly reduced by preincubation with PQQ. Moreover, in whole animal models of damage due to cardiac ischemia and reperfusion, PQQ results in less cardiac damage, higher left ventricle pressures, and fewer ventricular fibrillation episodes, if given i.p. 30 minutes before occlusion.⁶¹ In rodent models of cardiac ischemia, PQQ at doses ranging from 5-20 mg/kg administered i.p. was inversely related to infarct size. In the same tests, PQQ was superior to metoprolol in protecting mitochondria from ischemia/reperfusion oxidative damage.⁴³

Side Effects and Toxicity

Safety studies for PQQ in humans have been conducted in preparation for several human use patents.^{63,64} PQQ was administered at 20 or 60 mg/day for four weeks to two groups (10 each) of healthy adults given either a PQQ supplement or a placebo. These studies were double-blinded and conducted at two different

commercial drug-testing facilities: the New Drug Clinical Center, Fukuhara Clinic, Eniwa, Hokkaido, Japan and Cronova Co., Ltd., Suminoeku, Osaka, Japan. No adverse effects were observed in standard clinical tests at either dose (e.g., glucose, triglycerides, and various lipoprotein fractions). Functional tests for liver toxicity were also normal (e.g., aspartate aminotransferase and serum glutamic oxaloacetic transaminase). At 60 mg PQQ daily, the amounts of urinary N-acetyl- β -(D)-glucosaminidase activity were also within the normal range. N-acetyl-glucosaminidase is a renal hydrolytic enzyme located primarily in the lysosomal fraction of the renal tubular cell. Abnormal changes in renal tubular function or damage results in its elevation in urine.⁶⁵

Single-dose oral toxicity tests in rats were performed in compliance with Good Laboratory Practice (GLP). The single-dose oral toxicity tests indicated the approximate lethal dose of PQQ is less than 1,000 mg/kg body weight of rats, but higher than 500 mg/kg.

Post-mortem pathological examinations of test rats suggest the kidney as the principal target organ for acute effects of PQQ. In part, this is a validation of an earlier published toxicology study⁶⁶ in which PQQ was administered intraperitoneally to rats at a dose of 11-12 mg/kg body weight. Signs of renal tubular damage and inflammation were observed. When lower doses were used, however, no treatment effects or obvious pathological signs were observed. Likewise, in a 90-day repeated dose study in which PQQ was administered to rats by oral gavage (3, 20, or 100 mg PQQ/kg body weight) no adverse effects were observed. Moreover, at oral dosage levels from 250-2,000 mg PQQ/kg in mice, an examination for micronucleus induction in red blood cells showed no effects. Lastly, the results from a battery of genotoxicity tests *in vitro* (the Ames, micronucleus, and chromosomal aberration tests) were negative, i.e., PQQ did not cause clastogenic toxicity (chromosome breaks, rearrangements and changes in chromosomal number).

In summary, these observations taken together suggest there is no evidence of acute side effects or overt toxicity from consuming PQQ in amounts up to 60 mg per day for humans or several hundred mg per kg of diet fed to animals.

Dosage

Regarding typical exposures of free PQQ, as noted above, the amount for humans is estimated to vary from 100-400 μ g daily,^{11,25,28,67} about the same as the daily nutritional recommendations for biotin and folic acid, respectively. However, PQQ easily forms condensation products upon interaction with amino acids,²⁶ complicating the precision of such estimates. The primary condensation products are imidazolopyrroloquinoline (IPQ) and imidazolopyrroloquinoline derivatives with attached amino acid side chains as part of the chemical structure. For example, only about 15 percent of the PQQ is present in free form in biological fluids such as human milk, while 85 percent is present as IPQ and derivatives.²⁶ Thus, it is not unreasonable to assume that for humans the total exposure to PQQ derivatives may be as much as 1-2 mg per day. This amount is in the range that clearly influences optimization of growth and health in animal models.⁸ In the case of human milk, PQQ amounts to 1-2 μ g PQQ/g of milk solid, which is also similar to the PQQ concentrations reported for bovine milk.²⁶ It is important to note that PQQ appears readily absorbed. Smidt et al⁶⁸ determined that the apparent absorption of an oral dose of ¹⁴C-PQQ ranges from 20-80 percent when administered to adult mice in the fed state. The percentages were estimated from the amount of radioactivity present in urine and tissues 24 hours after administration.

Conclusions

The observation that increased mitochondrial genesis and antioxidant functions may be healthful features of PQQ supplementation opens the doors for both therapeutic applications and possible use as an ergogenic aid. Having normal mitochondrial function is essential to a broad range of health and disease relationships; thus, the need for continuing research that examines the efficacy and use of PQQ is compelling. PQQ derivatives are widely distributed in tissues and biological fluids at concentrations that may be sustained by typical dietary exposures. Given the range of functions and apparent survival benefits (e.g., improved reproductive performance), it is reasonable to suggest that PQQ may play a fundamental role in metabolism.

Acknowledgements

This article was supported in part by funds from the Mitsubishi Gas Chemical Co., the National Institutes of Health (NIH), and the Center for Health-Related Research, UC Davis.

References

- Hauge JG. Glucose dehydrogenase of *Bacterium anitratum*: an enzyme with a novel prosthetic group. *J Biol Chem* 1964;239:3630-3639.
- Anthony C. Pyrroloquinoline quinone (PQQ) and quinoprotein enzymes. *Antioxid Redox Signal* 2001;3:757-774.
- Anthony C, Williams P. The structure and mechanism of methanol dehydrogenase. *Biochim Biophys Acta* 2003;1647:18-23.
- Goodwin PM, Anthony C. The biochemistry, physiology and genetics of PQQ and PQQ-containing enzymes. *Adv Microb Physiol* 1998;40:1-80.
- Forrest HS, Salisbury SA, Sperl G. Crystallization of a derivative of a new coenzyme, methoxatin. *Biochim Biophys Acta* 1981;676:226-229.
- Forrest HS, Salisbury SA, Kilty CG. A mechanism for the enzymatic oxidation of methanol involving methoxatin. *Biochem Biophys Res Commun* 1980;97:248-251.
- Duine JA. Cofactor diversity in biological oxidations: implications and applications. *Chem Rec* 2001;1:74-83.
- Stites TE, Mitchell AE, Rucker RB. Physiological importance of quinoenzymes and the O-quinone family of cofactors. *J Nutr* 2000;130:719-727.
- Paz MA, Martin P, Fluckiger R, et al. The catalysis of redox cycling by pyrroloquinoline quinone (PQQ), PQQ derivatives, and isomers and the specificity of inhibitors. *Anal Biochem* 1996;238:145-149.
- Bishop A, Paz MA, Gallop PM, Karnovsky ML. Inhibition of redox cycling of methoxatin (PQQ), and of superoxide release by phagocytic white cells. *Free Radic Biol Med* 1995;18:617-620.
- Fluckiger R, Paz MA, Gallop PM. Redox-cycling detection of dialyzable pyrroloquinoline quinone and quinoproteins. *Methods Enzymol* 1995;258:140-149.
- Karnovsky ML, Bishop A, Camerero VC, et al. Aspects of the release of superoxide by leukocytes, and a means by which this is switched off. *Environ Health Perspect* 1994;102:43-44.
- Paz MA, Fluckiger R, Gallop PM. Redox-cycling is a property of PQQ but not of ascorbate. *FEBS Lett* 1990;264:283-284.
- Katz E, Willner I. A biofuel cell with electrochemically switchable and tunable power output. *J Am Chem Soc* 2003;125:6803-6813.
- Mukherjee J, Lumibao CY, Kirchoff JR. Application of a thiol-specific electrocatalytic electrode for real-time amperometric monitoring of enzymatic hydrolysis. *Analyst* 2009;134:582-586.
- Willner I, Baron R, Willner B. Integrated nanoparticle-biomolecule systems for biosensing and bioelectronics. *Biosens Bioelectron* 2007;22:1841-1852.
- Szeponik J, Möller B, Pfeiffer D, et al. Ultrasensitive bienzyme sensor for adrenaline. *Biosens Bioelectron* 1997;12:947-952.
- Puehringer S, Metlitzky M, Schwarzenbacher R. The pyrroloquinoline quinone biosynthesis pathway revisited: a structural approach. *BMC Biochem* 2008;9:8.
- Magnusson OT, RoseFigura JM, Toyama H, et al. Pyrroloquinoline quinone biogenesis: characterization of PqqC and its H84N and H84A active site variants. *Biochemistry* 2007;46:7174-7186.
- Magnusson OT, Toyama H, Saeki M, et al. Quinone biogenesis: structure and mechanism of PqqC, the final catalyst in the production of pyrroloquinoline quinone. *Proc Natl Acad Sci U S A* 2004;101:7913-7918.
- Krueger FR, Werther W, Kissel J, Schmid ER. Assignment of quinone derivatives as the main compound class composing 'interstellar' grains based on both polarity ions detected by the 'Cometary and Interstellar Dust Analyser' (CIDA) onboard the spacecraft STARDUST. *Rapid Commun Mass Spectrom* 2004;18:103-111.
- Kissel J, Krueger FR, Silén J, Clark BC. The Cometary and Interstellar Dust Analyser at comet 81P/Wild 2. *Science* 2004;304:1774-1776.
- Choi O, Kim J, Kim JG, et al. Pyrroloquinoline quinone is a plant growth promotion factor produced by *Pseudomonas fluorescens* B16. *Plant Physiol* 2008;146:657-668.
- Rodriguez H, Gonzalez T, Selman G. Expression of a mineral phosphate solubilizing gene from *Erwinia herbicola* in two rhizobacterial strains. *J Biotechnol* 2001;84:155-161.
- Pierpoint WS. PQQ in plants. *Trends Biochem Sci* 1990;15:299.
- Mitchell AE, Jones AD, Mercer RS, Rucker RB. Characterization of pyrroloquinoline quinone amino acid derivatives by electrospray ionization mass spectrometry and detection in human milk. *Anal Biochem* 1999;269:317-325.

27. Kumazawa T, Seno H, Suzuki O. Failure to verify high levels of pyrroloquinoline quinone in eggs and skim milk. *Biochem Biophys Res Commun* 1993;193:1-5.
28. Kumazawa T, Sato K, Seno H, et al. Levels of pyrroloquinoline quinone in various foods. *Biochem J* 1995;307:331-333.
29. Matsushita K, Arents JC, Bader R, et al. *Escherichia coli* is unable to produce pyrroloquinoline quinone (PQQ). *Microbiology* 1997;143:3149-3156.
30. Steinberg FM, Gershwin ME, Rucker RB. Dietary pyrroloquinoline quinone: growth and immune response in BALB/c mice. *J Nutr* 1994;124:744-753.
31. Steinberg F, Stites TE, Anderson P, et al. Pyrroloquinoline quinone improves growth and reproductive performance in mice fed chemically defined diets. *Exp Biol Med* (Maywood) 2003;228:160-166.
32. Kasahara T, Kato T. Nutritional biochemistry: a new redox-cofactor vitamin for mammals. *Nature* 2003;422:832.
33. Rucker R, Storms D, Sheets A, et al. Biochemistry: is pyrroloquinoline quinone a vitamin? *Nature* 2005;433:E10-E11;discussion E11-E12.
34. Felton LM, Anthony C. Biochemistry: role of PQQ as a mammalian enzyme cofactor? *Nature* 2005;433:E10;discussion E11-E12.
35. Sato K, Toriyama M. Effect of pyrroloquinoline quinone (PQQ) on melanogenic protein expression in murine B16 melanoma. *J Dermatol Sci* 2009;53:140-145.
36. Hirakawa A, Shimizu K, Fukumitsu H, Furukawa S. Pyrroloquinoline quinone attenuates iNOS gene expression in the injured spinal cord. *Biochem Biophys Res Commun* 2009;378:308-312.
37. Nunome K, Miyazaki S, Nakano M, et al. Pyrroloquinoline quinone prevents oxidative stress-induced neuronal death probably through changes in oxidative status of DJ-1. *Biol Pharm Bull* 2008;31:1321-1326.
38. Kumazawa T, Hiwasa T, Takiguchi M, et al. Activation of ras signaling pathways by pyrroloquinoline quinone in NIH3T3 mouse fibroblasts. *Int J Mol Med* 2007;19:765-770.
39. Chohanadisai W, Bauerly K, Tchapanian E, Rucker RB. Pyrroloquinoline quinone (PQQ) stimulates mitochondrial biogenesis. *FASEB J* 2007;21:854.
40. Ouchi A, Nakano M, Nagaoka S, Mukai K. Kinetic study of the antioxidant activity of pyrroloquinolinequinol (PQQH(2)), a reduced form of pyrroloquinolinequinone in micellar solution. *J Agric Food Chem* 2009;57:450-456.
41. Stites T, Storms D, Bauerly K, et al. Pyrroloquinoline quinone modulates mitochondrial quantity and function in mice. *J Nutr* 2006;136:390-396.
42. Bauerly KA, Storms DH, Harris CB, et al. Pyrroloquinoline quinone nutritional status alters lysine metabolism and modulates mitochondrial DNA content in the mouse and rat. *Biochim Biophys Acta* 2006;1760:1741-1748.
43. Zhu BQ, Simonis U, Cecchini G, et al. Comparison of pyrroloquinoline quinone and/or metoprolol on myocardial infarct size and mitochondrial damage in a rat model of ischemia/reperfusion injury. *J Cardiovasc Pharmacol Ther* 2006;11:119-128.
44. Debray FG, Lambert M, Mitchell GA. Disorders of mitochondrial function. *Curr Opin Pediatr* 2008;20:471-482.
45. Puigserver P. Tissue-specific regulation of metabolic pathways through the transcriptional coactivator PGC1-alpha. *Int J Obes (Lond)* 2005;29:S5-S9.
46. Muoio DM, Koves TR. Skeletal muscle adaptation to fatty acid depends on coordinated actions of the PPARs and PGC-1alpha: implications for metabolic disease. *Appl Physiol Nutr Metab* 2007;32:874-883.
47. Tsuchida T, Yasuyama T, Higuchi K, et al. The protective effect of pyrroloquinoline quinone and its derivatives against carbon tetrachloride-induced liver injury of rats. *J Gastroenterol Hepatol* 1993;8:342-347.
48. Urakami T, Yoshida C, Akaike T, et al. Synthesis of monoesters of pyrroloquinoline quinone and imidazopyrroloquinoline, and radical scavenging activities using electron spin resonance *in vitro* and pharmacological activity *in vivo*. *J Nutr Sci Vitaminol (Tokyo)* 1997;43:19-33.
49. Hamagishi Y, Murata S, Kamei H, et al. New biological properties of pyrroloquinoline quinone and its related compounds: inhibition of chemiluminescence, lipid peroxidation and rat paw edema. *J Pharmacol Exp Ther* 1990;255:980-985.
50. Naito Y, Kumazawa T, Kino I, Suzuki O. Effects of pyrroloquinoline quinone (PQQ) and PQQ-oxazole on DNA synthesis of cultured human fibroblasts. *Life Sci* 1993;52:1909-1915.
51. Aizenman E, Hartnett KA, Zhong C, et al. Interaction of the putative essential nutrient pyrroloquinoline quinone with the N-methyl-D-aspartate receptor redox modulatory site. *J Neurosci* 1992;12:2362-2369.
52. Zhang Y, Feustel PJ, Kimelberg HK. Neuroprotection by pyrroloquinoline quinone (PQQ) in reversible middle cerebral artery occlusion in the adult rat. *Brain Res* 2006;1094:200-206.
53. Zhang Y, Rosenberg PA. The essential nutrient pyrroloquinoline quinone may act as a neuroprotectant by suppressing peroxynitrite formation. *Eur J Neurosci* 2002;16:1015-1024.

54. Jensen FE, Gardner GJ, Williams AP, et al. The putative essential nutrient pyrroloquinoline quinone is neuroprotective in a rodent model of hypoxic/ischemic brain injury. *Neuroscience* 1994;62:399-406.
55. Sanchez RM, Wang C, Gardner G, et al. Novel role for the NMDA receptor redox modulatory site in the pathophysiology of seizures. *J Neurosci* 2000;20:2409-2417.
56. Scanlon JM, Aizenman E, Reynolds IJ. Effects of pyrroloquinoline quinone on glutamate-induced production of reactive oxygen species in neurons. *Eur J Pharmacol* 1997;326:67-74.
57. Aizenman E, Jensen FE, Gallop PM, et al. Further evidence that pyrroloquinoline quinone interacts with the N-methyl-D-aspartate receptor redox site in rat cortical neurons *in vitro*. *Neurosci Lett* 1994;168:189-192.
58. Zhang P, Xu Y, Sun J, et al. Protection of pyrroloquinoline quinone against methylmercury-induced neurotoxicity via reducing oxidative stress. *Free Radic Res* 2009;43:224-233.
59. Murase K, Hattori A, Kohno M, Hayashi K. Stimulation of nerve growth factor synthesis/secretion in mouse astroglial cells by coenzymes. *Biochem Mol Biol Int* 1993;30:615-621.
60. Ohwada K, Takeda H, Yamazaki M, et al. Pyrroloquinoline quinone (PQQ) prevents cognitive deficit caused by oxidative stress in rats. *J Clin Biochem Nutr* 2008;42:29-34.
61. Tao R, Karliner JS, Simonis U, et al. Pyrroloquinoline quinone preserves mitochondrial function and prevents oxidative injury in adult rat cardiac myocytes. *Biochem Biophys Res Commun* 2007;363:257-262.
62. Zhu BQ, Zhou HZ, Teerlink JR, Karliner JS. Pyrroloquinoline quinone (PQQ) decreases myocardial infarct size and improves cardiac function in rat models of ischemia and ischemia/reperfusion. *Cardiovasc Drugs Ther* 2004;18:421-431.
63. Tsuji T, Yamaguchi K, Kondo K, Urakami T. Nerve growth factor production accelerators and compositions for preventing or treating neuronal degeneration. US Patent 5846977; 1998.
64. Urakami T. Process for the preparation of pyrroloquinoline quinone. US Patent 5344768; September 6, 1994.
65. D'Amico G, Bazzi C. Urinary protein and enzyme excretion as markers of tubular damage. *Curr Opin Nephrol Hypertens* 2003;12:639-643.
66. Watanabe A, Hobara N, Ohsawa T, et al. Nephrotoxicity of pyrroloquinoline quinone in rats. *Hiroshima J Med Sci* 1989;38:49-51.
67. Fluckiger R, Paz M, Mah J, et al. Characterization of the glycine-dependent redox-cycling activity in animal fluids and tissues using specific inhibitors and activators: evidence for presence of PQQ. *Biochem Biophys Res Commun* 1993;196:61-68.
68. Smidt CR, Unkefer CJ, Houck DR, Rucker RB. Intestinal absorption and tissue distribution of [¹⁴C] pyrroloquinoline quinone in mice. *Proc Soc Exp Biol Med* 1991;197:27-31.

Corrections

Altern Med Rev 2009;14(2):143.

Figure 2. Meriva should read Meriva curcuminoids, curcumin should read curcuminoids.

Paragraph 3: "One small unpublished..."

Should read: "One small unpublished, single-dose trial demonstrated 450 mg of Meriva curcuminoids complexed with phosphatidylcholine was absorbed as efficiently as 4 g unbound *Curcuma longa* (95% curcumin), reflecting a significant increase in bioavailability for Meriva complex (Figure 2).15